

### **III. MISSION OVERVIEW AND SUMMARIES OF EXPERIMENTS**

#### **EFFECTS OF MICROGRAVITY ON RAT MUSCLE**

**D.A. Riley**

**Department of Anatomy and Cellular Biology  
Medical College of Wisconsin  
Milwaukee, Wisconsin 53225**

It is well known that humans exposed to long term spaceflight experience undesirable progressive muscle weakness and increased fatigability. This problem has prompted the implementation of inflight exercise programs because most investigators believe that the major cause of diminished muscle performance is a combination of disuse and decreased workload. Inflight exercise has improved muscle health, but deficits have persisted, indicating that either the regimens utilized were suboptimal or there existed additional debilitating factors which were not remedied by exercise.

Clarification of this question requires an improved understanding of the cellular and molecular basis of spaceflight-induced muscle deterioration. To this end, multiple investigations have been performed on the muscles from rats orbited 5-22 days in Cosmos biosatellites and Spacelab-3 (2,4,5,8,10-14,16,18,19, 21-23,25,27,28).

The eight Cosmos 1887 investigations that follow examined the structural and biochemical changes in skeletal and cardiac muscles of rats exposed to microgravity for 12.5 days and returned to terrestrial gravity 2.3 days before tissues were collected. Even though interpretation of these results was complicated by the combination of inflight and postflight induced alterations, the consensus is that there is marked heterogeneity in both the degree and type of responses from the whole muscle level down to the molecular level. Consistent with previous reports, red (oxidative) antigravity muscles (such as soleus, adductor longus, and vastus intermedius) were more atrophic than white (glycolytic) non-antigravity muscles (extensor digitorum longus, tibialis anterior, vastus lateralis and medialis) (2,4,5,13,14,16,17, 19,23,25,27). To some extent, fiber type composition predicts the degree of muscle atrophy. The most affected muscles were composed primarily of slow oxidative fibers whereas fast oxidative glycolytic and fast glycolytic fibers predominated in the least affected muscles. However, at another level of complexity, fibers of a given type located deep in a muscle (closer to the bone) exhibited greater atrophy than fibers of the same type located superficially in the same muscle (7). More limited in occurrence than muscle atrophy was the degeneration of muscle fibers, nerves, and micro-vessels which only involved the adductor longus and soleus muscles (24). Within the adductor longus, the pathology affected the mid-belly or endplate region more extensively than the ends or myotendinous zones.

Generally speaking, muscle enzyme properties shifted from slow oxidative toward fast glycolytic, but microheterogeneity was evident. During atrophy, contractile proteins, especially slow myosin, were preferentially lost relative to cytoplasmic non-contractile proteins (1,7,28). Increased ubiquitination of contractile proteins was observed (24). This process is postulated to promote selective protein degradation (26). Within slow oxidative fibers, slow myosin was replaced by fast myosin resulting in hybrid slow/fast fibers with elevated myosin ATPase activity (1,7). The shifts in cytoplasmic enzymes were not uniform. In soleus muscles, oxidative enzymes (citrate synthase and malate, beta-hydroxyacyl CoA, and

succinate dehydrogenases) decreased in step with muscle atrophy so that their concentrations were minimally altered (15). In contrast, glycogenolytic enzymes (glycogen phosphorylase, glycerophosphate dehydrogenase, lactate dehydrogenase, and pyruvate kinase) were not lost and their amounts per fiber increased during atrophy. In contrast to soleus, individual muscle fibers of the tibialis anterior showed elevated oxidative enzymes in atrophic fibers whereas glycogenolytic enzymes minimally changed in concentration (15). Mitochondrial enzyme changes were not uniform across the diameter of a muscle fiber. Mitochondria and their associated enzyme activities were essentially unchanged centrally but showed a marked decrease in the peripheral subsarcolemmal population (7,24). The mechanism by which proteins are turned over selectively and enzyme levels are altered within specific regions of atrophic muscle fibers are not understood.

The absence of detectable shifts in alpha-actin and cytochrome c mRNA levels may have resulted from the postflight muscle contractile activity because messenger levels can easily be altered within two days (29). Other parameters, such as atrophy, do not change markedly in two days and therefore, more accurately reflect microgravity induced changes. Distinguishing between microgravity induced and postflight readaptation induced alterations is a primary objective for the proposed studies in which muscles are harvested immediately (within 4 to 9 hours) upon landing (Cosmos July 1989, SLS-1) or inflight (SLS-2).

The muscle atrophy associated with spaceflight is most likely due to disuse and reduced workload. The observed decrease in mature collagen cross-links, collagen concentration and DNA concentration in the patellar tendons of flight rats is consistent with lowered workload (30). However, the Achilles tendon, serving the gastrocnemius/soleus/plantaris complex, was unchanged in the same animals (3). Again, these findings illustrate the diversity of skeletal muscle tissue responses. Activity and workload can be assessed by instrumenting muscles with electromyographic electrodes and tendon tension transducers. Systematic factors may have also contributed to atrophy. The suspected lower levels of growth hormone would have had a general catabolic effect on a skeletal muscle and elevated glucocorticoids, indicated by adrenal cortex hypertrophy, would have specifically induced fast fiber atrophy (9).

Some cardiac and skeletal muscle fibers were necrotic (20,24). The motor innervation of skeletal muscle was partially degenerated (24). Microcirculatory vessels were disrupted in skeletal muscle and abnormal in cardiac muscle (20,24). Repair of necrotic muscle fibers, motor axons, and blood vessels is dependent upon the effectiveness of complex processes of regeneration. Ground-based evidence predicts that regeneration may be compromised in space because of reduced active muscle tension and exposure of the dividing stem cells to damaging cosmic radiation (3,6,13). The ability of tissues to regenerate effectively during spaceflight is important to address in future missions.

Collectively, the muscle investigations of Cosmos 1887 clearly illustrate the wide diversity of muscle tissue responses to spaceflight. Judging from the summary report of this mission, heterogeneity of responses is not unique to muscle tissue. Elucidating the mechanism underlying this heterogeneity holds the key to explaining adaptation of the organism to prolonged spaceflight.

## REFERENCES

1. Baldwin, K.M. and Herrick, R.E. Effect of Zero Gravity on Myosin Isoform Expression in Rodent Skeletal Muscle. In Final Reports of U.S. Experiments Flown on the Soviet Biosatellite Cosmos 1887. (Connolly, J.P., Grindeland, R.E., and Ballard, R.W., eds). This report.
2. Baranski, S., Baranska, W., Marciniak, M., Ilyina-Kakueva, E.I. 1979. Ultrasonic ("Ultrastructural") investigations of the soleus muscle after space flight on the Biosputnik 936. *Aviat. Space Environ. Med.* 50:930-934.
3. Benton, E.V. Peterson, D.D., Marennny, A.M., Popov, V.I. 1978. HZE particle radiation studies aboard Kosmos 782. *Health Physics.* 35:643-648.
4. Castleman K.R et al. 1978. Spaceflight effects on muscle fibers: Final Reports of U.S. Experiments Flown on Soviet Satellite Cosmos 936. Rosenzweig, S.N., Souza, K.A., Eds. NASA Technical Memorandum 78526. pp. 224-289.
5. Castleman, K.R., Chui, L.A., Van Der Meulen, J.P. 1981. Automatic analysis of muscle fibers from rats subjected to spaceflight. NASA Technical Memorandum 81289. pp. 267-277.
6. Denny-Brown, D. 1951. The influence of tension and innervation on the regeneration of skeletal muscle. *J. Neuropath. Exp. Neurol.* 10:94-95.
7. Edgerton, V.R., Martin, T.P., Roy R.R., Marini, J.F., Leger, J.J. Metabolic and Morphological Properties of Muscle Fibers after Spaceflight. In Final Reports of U.S. Experiments Flown on the Soviet Biosatellite Cosmos 1887. (Connolly, J.P., Grindeland, R.E., and Ballard, R.W., eds). This report.
8. Gazenko, O.G., Genin, A.M. Il'in, Y.A, Portugalov, V.V., Serova, L.V., Tigranyan, R.A. 1978. Principal results of experiment with mammals onboard the Kosmos-782 biosatellite. *Kosmich. Biol. Aviakosmisch. Med.* 6:43-49.
9. Grindeland, R.E., Popova, I., Vasques, M. Effects of Microgravity on Rat Body and Adrenal Weights and Plasma Constituents. In Final Reports of U.S. Experiments Flown on the Soviet Biosatellite Cosmos 1887. (Connolly, J.P., Grindeland, R.E., and Ballard, R.W., eds). This report.
10. Ilyin, E.A. 1983. Investigations on biosatellites of the Cosmos series. *Aviat. Space Environ. Med. Suppl. I.* 54:S9-S13.
11. Ilyina-Kakueva, E.I. 1987. Investigation of rat skeletal muscles following short-term spaceflight aboard Cosmos-1667 biosatellite. *Kosmich. Biol. Aviakosmich. Med.* 21:31-35.
12. Ilyina-Kakueva, E.I. Portugalov, V.V. 1981. Structural changes in the soleus muscle of rats flown aboard the Cosmos series of biosatellites and submitted to hypokinesia. *Kosmich. Biol. Aviakosmich. Med.* 15:37-40.

13. Ilyina-Kakuyeva, E.I. Portugalov, V.V. 1977. Combined effect of space flight and radiation on skeletal muscles of rats. *Aviat. Space Environ. Med.* 48:115-119.
14. Ilyina-Kakueva, E.I. Portugalov, V.V., Krivenkova, N.P. 1976. Space flight effects on the skeletal muscle of rats. *Aviat. Space Environ. Med.* 47:700-703.
15. Lowry, O.H. McDougal, D.B., Chi, M.-Y., Pusateri, M.E., Carter, J.G., Manchester, J.K. Effect of Microgravity on 1) Metabolic enzymes of Type 1 and Type 2 Muscle Fibers and on 2) Metabolic Enzymes, Neurotransmitter Amino Acids and Neurotransmitter Associated Enzymes in Motor and Somatosensory Cerebral Cortex. In Final Reports of U.S. Experiments Flown on the Soviet Biosatellite Cosmos 1887. (Connolly, J.P., Grindeland, R.E., and Ballard, R.W., eds). This report.
16. Martin, T.P. Edgerton, V.R., Grindeland, R.E. 1988. Influence of spaceflight on rat skeletal muscle. *J. Appl Physiol.* 65:2318-2325.
17. Musacchia, X.J., Steffen, J.M., Fell, R.D. Skeletal Muscle Atrophy in Response to 14 Days of Weightlessness. In Final Reports of U.S. Experiments Flown on the Soviet Biosatellite Cosmos 1887. (Connolly, J.P., Grindeland, R.E., and Ballard, R.W., eds). This report.
18. Oganov, V.S., Skuratova, S.A., Potapov, A.N. Shirvinskaya, K., Marusich, M. 1982. Physiological mechanisms of adaptation of rat skeletal muscles to weightlessness and similar functional requirements. *The Physiologist.* 23:S16-S21.
19. Oganov, V.S., Rakhmanov, A.S., Skwatova, S.A., Shirvinskaya, Magedov, V.S. 1985. Functions of skeletal muscles of rats and monkeys after 5-day spaceflight on Cosmos 1514. *Proc. 2nd Int. Conf. Space Physiol., Toulouse, France.*
20. Philpott, D.E., Mednieks, M.I., Papova, I.A., Serova, I.V., Kato, K. Stevenson, J., Miquel, J. Sapp, W. Morphological and Biochemical Examination of Heart Tissue.
21. Portugalov, V.V., Ilyina-Kakueva, E.I. Prolonged spaceflight and hypokinesia. *Aerospace Med.* 44:764-768.
22. Portugalov, V.V., Ilyina-Kakueva, E.I. Starostin, V.I., Rokhlenko, D.K., Savik, Z.F. 1971. Morphological and cytochemical studies of hypokinetic effects. *Aerospace Med.* 42:1041-1049.
23. Rapcsak, M., Oganov, V.S., Szoor, A., Skuratova, S.A., Szilagyi, R. 1983. Effect of weightlessness on the function of rat skeletal muscles on the biosatellite Cosmos-1129. *Acta Physiol. Hung.* 62:225-228.
24. Riley, D.A., Ilyina-Kakueva, E.I., Ellis, S., Bain, J.L.W., Sedlak, R.F., Slocum, G.R. Morphological, Histochemical and Biochemical Investigation of Microgravity-induced Nerve and Muscle Breakdown. In Final Reports of U.S. Experiments Flown on the Soviet Biosatellite Cosmos 1887. (Connolly, J.P., Grindeland, R.E., and Ballard, R.W., eds). This report.

25. Riley, D.A., Ellis, S., Slocum, G.R., Satyanarayana, T., Bain, J.L.W., Sedlak, F.R. 1987. Hypogravity-induced atrophy of rat soleus and extensor digitorum longus muscles. *Muscle & Nerve*. 10:560-568.
26. Riley, D.A., Bain, J.L.W., Ellis, S., Haas, A.L. 1988. Quantification and immunocytochemical localization of ubiquitin conjugates within rat red and white skeletal muscles. *J. Histochem. Cytochem.* 36:621-632.
27. Rokhlenko, K.D., Savik, Z.F. 1981. Effect of space flight factors on ultrastructure of skeletal muscles. *Kosmich. Biol. Aviakosmich. Med.* 1:72-77.
28. Takacs, O., Rancsak, M., Szoor, A., Oganov, V.S., Szilagyi, T., Oganessian, S.S., Guba, F. 1983. Effect of weightlessness on myofibrillar proteins of rat skeletal muscles with different functions in experiment of biosatellite Cosmos-1129. *Acta Physiol. Hung.* 62:228-233.
29. Thomason, D.B., Morrison, P.R., Booth, F.W. Messenger RNA Levels in Skeletal Muscle. In *Final Reports of U.S. Experiments Flown on the Soviet Biosatellite Cosmos 1887*. (Connolly, J.P., Grindeland, R.E., and Ballard, R.W., eds). This report.
30. Vailas, A.C., Zernicke, R.F., Grindeland, R.E. Biomechanical, Biochemical and Morphological Alterations of Muscle and Dense, Fibrous Connective Tissues during 14 Days of Spaceflight. In *Final Reports of U.S. Experiments Flown on the Soviet Biosatellite Cosmos 1887*. (Connolly, J.P., Grindeland, R.E., and Ballard, R.W., eds). This report.